arXiv:1408.4871v1 [math.AT] 21 Aug 2014

CERTAIN NUMBERS ON THE GROUPS OF SELF-HOMOTOPY EQUIVALENCES

Ho Won Choi and Kee Young Lee

Abstract. For a connected based space X, let $[X, X]$ be the set of all based homotopy classes of base point preserving self map of X and let $\mathcal{E}(X)$ be the group of self-homotopy equivalences of X. We denote by $\mathcal{A}_\sharp^k(X)$ the set of homotopy classes of self-maps of X that induce an automorphism of $\pi_i(X)$ for $i = 0, 1, \cdots, k$. That is, $[f] \in \mathcal{A}_\sharp^k(X)$ if and only if $\pi_i(f) : \pi_i(X) \to \pi_i(X)$ is an isomorphism for $i = 0, 1, \cdots, k$. Then, $\mathcal{E}(X) \subseteq \mathcal{A}_\sharp^k(X) \subseteq [X, X]$ for a nonnegative integer k. Moreover, for a connected CW-complex X, we have $\mathcal{E}(X) = \mathcal{A}_\sharp(X)$. In this paper, we study the properties of $\mathcal{A}_\sharp^k(X)$ and discuss the conditions under which $\mathcal{E}(X) = \mathcal{A}_\sharp^k(X)$ and the minimum value of such k. Furthermore, we determine the value of k for various spaces, including spheres, products of spaces, and Moore spaces.

1. Introduction

For a connected space X, we denote by $[X, X]$ the set of all based homotopy classes of self maps of X. Then, $[X, X]$ is a monoid with multiplication given by the composition of homotopy classes. Let $\mathcal{E}(X)$ be the set of self-homotopy equivalences of X. Then, $\mathcal{E}(X)$ is a group with the operation given by the composition of homotopy classes. $\mathcal{E}(X)$ has been studied extensively by various authors, including Arkowitz[1], Maruyama[4], Lee[12], Oka[13], Rutter[15], Sawashita[16], and Sieradski[17]. Several subgroups of $\mathcal{E}(X)$ have also been studied. One of these is $\mathcal{E}_\sharp^n(X)$ which consists of the self-homotopy equivalences that induce the identity homomorphism on homotopy groups $\pi_i(X)$, for $i = 1, 2, \cdots, n$. That is, $\mathcal{E}_\sharp^n(X) = \{f \in \mathcal{E}(X) \mid \pi_i(f) = id_{\pi_i(X)} \text{ on } \pi_i(X) \text{ for } i = 0, \cdots, n\}$, where $\pi_i(f) : \pi_i(X) \to \pi_i(X)$ is the induced homomorphism of a map $f : X \to X$ and $n = \infty$ is available. When $n = \infty$, we simply denote $\mathcal{E}_\sharp^\infty(X)$ as $\mathcal{E}_\sharp(X)$. It is well known that $\mathcal{E}(S^n) = Z_2$ and $\mathcal{E}_\sharp(S^n) = 1$, where S^n is the n-dimensional sphere. Moreover, $\mathcal{E}_\sharp^n(X)$ is a subgroup of $\mathcal{E}(X)$ and, if $n \leq m$, $\mathcal{E}_\sharp^m(X) \subseteq \mathcal{E}_\sharp^n(X)$. Thus, we have the following chain by inclusion:

$$\mathcal{E}_\sharp(X) \subseteq \cdots \subseteq \mathcal{E}_\sharp^n(X) \subseteq \cdots \subseteq \mathcal{E}_\sharp^1(X) \subseteq \mathcal{E}(X).$$

2000 *Mathematics Subject Classification.* Primary 55P10, 555Q05, 55Q52.

Key words and phrases. homotopy equivalence, self closeness number, k-self equivalence.

This work was supported by a Korea University Grant.

We denote by $\mathcal{A}_\sharp^k(X)$ the set of homotopy classes of self-maps of X that induce an automorphism of $\pi_i(X)$ for $i = 0, 1, \cdots, k$. That is, $[f] \in \mathcal{A}_\sharp^k(X)$ if and only if $\pi_i(f) : \pi_i(X) \to \pi_i(X)$ is an isomorphism for $i = 0, 1, \cdots, k$. Then, for any nonnegative integer k, $\mathcal{A}_\sharp^k(X)$ is a submonoid of $[X, X]$ and always contains $\mathcal{E}(X)$. $n = \infty$ is available and, if $n = \infty$, we simply denote $\mathcal{A}_\sharp^\infty(X)$ as $\mathcal{A}_\sharp(X)$. If $k < n$, then $\mathcal{A}_\sharp^n(X) \subseteq \mathcal{A}_\sharp^k(X)$; thus, we have the following chain by inclusion:

$$\mathcal{E}(X) \subseteq \mathcal{A}_\sharp(X) \subseteq ... \subseteq \mathcal{A}_\sharp^1(X) \subseteq \mathcal{A}_\sharp^0(X) = [X, X].$$

For any connected CW-complex X, we have $\mathcal{A}_\sharp(X) = \mathcal{E}(X)$. These two facts give rise to the question of what is the largest $\mathcal{A}_\sharp^n(X)$ that is equal to $\mathcal{E}(X)$. In other words, what is the minimum k such that $\mathcal{E}(X) = \mathcal{A}_\sharp^k(X)$? We denote by $N\mathcal{E}(X)$ the least nonnegative integer k such that $\mathcal{E}(X) = \mathcal{A}_\sharp^k(X)$, and call it *the self-closeness number of* X. That is,

$$N\mathcal{E}(X) = min\{k \mid \mathcal{A}_\sharp^k(X) = \mathcal{E}(X) \ for \ k \geq 0\}.$$

In this paper, we investigate properties of the self-closeness number of a space including the homotopy invariance, the relation with the dimension of spaces and the relation with the n-connectivity. From the results of this investigation, we completely determine the value of k in several specific cases. Moreover, we give an equivalence relation on $\mathcal{A}_\sharp^k(X)$ and study the relations between the set of equivalence classes and $\mathcal{E}(X)$, $\mathcal{E}_\sharp^n(X)$, or the self-closeness numbers.

Throughout this paper, all topological spaces are based and have the based homotopy type of CW-complex. All maps and homotopies will preserve the base points. For the spaces X and Y, we denote by $[X, Y]$ the set of homotopy classes of maps from X to Y. We do not distinguish between the notation of a map $X \to Y$ and that of its homotopy class in $[X, Y]$.

2. Properties of $\mathcal{A}_\sharp^k(X)$ and the self-closeness number

We begin by identifying $\mathcal{A}_\sharp^k(X)$ as a submonoid of $[X, X]$.

Lemma 1. *$\mathcal{A}_\sharp^k(X)$ is a submonoid of $[X, X]$ for any nonnegative integer k.*

Proof. Let $f, g \in \mathcal{A}_\sharp^k(X)$. Then, $\pi_i(f), \pi_i(g) \in Aut(\pi_i(X))$ for $i = 0, 1, \cdots, k$, where $Aut(\pi_i(X))$ is the automorphism group of $\pi_i(X)$. Thus, $\pi_i(f \circ g) = \pi_i(f) \circ \pi_i(g) \in Aut(\pi_i(X))$ for $i = 0, 1, \cdots, k$. It follows that $f \circ g \in \mathcal{A}_\sharp^k(X)$. Clearly, $f \circ (g \circ h) = (f \circ g) \circ h$. Moreover, because $\pi_i(id_X) = id_{\pi_i(X)} \in Aut(\pi_i(X))$ for $i = 0, 1, \cdots, k$, $id_X \in \mathcal{A}_\sharp^k(X)$. □

In general, $\mathcal{E}(X) \neq \mathcal{A}_\sharp(X)$.

Example 1. Let X be a quasi-circle in the plane $\mathbb{R}^2$; that is,

$$X = \{(x, y)|\ y = sin(\frac{1}{x}), 0 < x \leq \frac{1}{\pi}\} \cup \{(0, y)| -2 \leq y \leq 1\}$$

$$\cup\{(x,-2)|0\leq x\leq\frac{1}{\pi}\}\cup\{(\frac{1}{\pi},y)|-2\leq y\leq 0\}.$$

Then, for each $x_0\in X$, the inclusion map $i:x_0\to X$ is a weak homotopy equivalence, but not a homotopy equivalence. Thus, the constant map $C:X\to X$ given by $C(x)=x_0$ is not a homotopy equivalence. Therefore, C does not belong to $\mathcal{E}(X)$. However, as $\pi_i(X)=0$ for all i, $\pi_i(C)\in Aut(\pi_i(X))$ for all i.

Whitehead theorem ensures that $\mathcal{E}(X)=\mathcal{A}_\sharp(X)$ provided that X is a CW-complex.

Lemma 2. *If X is a CW-complex, then $\mathcal{E}(X)=\mathcal{A}_\sharp(X)$.*

In general, $\mathcal{A}_\sharp^{k-1}(X)\neq\mathcal{A}_\sharp^k(X)$.

Example 2. Let $M(\mathbb{Z}_4,5)$ be a Moore space. Then, by the results in [6], $[M(\mathbb{Z}_4,5),M(\mathbb{Z}_4,5)]\cong\mathbb{Z}_4\oplus\mathbb{Z}_2$. Since $M(\mathbb{Z}_4,5)$ is 4-connected, $\pi_i(M(\mathbb{Z}_4,5))=0$ for $i=0,1,2,3,4$. Thus, $\mathcal{A}_\sharp^4(M(\mathbb{Z}_4,5))=[M(\mathbb{Z}_4,5),M(\mathbb{Z}_4,5)]\cong\mathbb{Z}_4\oplus\mathbb{Z}_2$ and, for $f\in[M(\mathbb{Z}_4,5),M(\mathbb{Z}_4,5)]$, $f\in\mathcal{A}_\sharp^5(M(\mathbb{Z}_4,5))$ if and only if $\pi_5(f)\in Aut(\pi_5(M(\mathbb{Z}_4,5)))$. Therefore, $\mathcal{A}_\sharp^5(M(\mathbb{Z}_4,5))\cong\mathbb{Z}_4^*\times\mathbb{Z}_2$, where $\mathbb{Z}_4^*$ is the automorphism group of $\mathbb{Z}_4$. Rutter [15] showed that $\mathcal{E}(M(\mathbb{Z}_4,5))\cong\mathbb{Z}_4^*\times\mathbb{Z}_2$. Consequently,

$$\mathcal{A}_\sharp^5(M(\mathbb{Z}_4,5))=\mathcal{E}(M(\mathbb{Z}_4,5))\cong\mathbb{Z}_4^*\times\mathbb{Z}_2\cong\mathbb{Z}_2\times\mathbb{Z}_2.$$

Therefore, we have

$$\mathcal{E}(M(\mathbb{Z}_4,5))=\mathcal{A}_\sharp^5(M(\mathbb{Z}_4,5))\subsetneq\mathcal{A}_\sharp^4(M(\mathbb{Z}_4,5))=[M(\mathbb{Z}_4,5),M(\mathbb{Z}_4,5)].$$

Example 2 ensures that the self-closeness number $N\mathcal{E}(M(\mathbb{Z}_4,5)$ is 5. We will discuss details of the more general Moore spaces $M(G,n)$ where G is an abelian group in Section 3.

The self-closeness number is a homotopy invariant.

Theorem 1. *Let X and Y be CW-complexes. If X and Y have the same homotopy type, then $N\mathcal{E}(X)=N\mathcal{E}(Y)$.*

Before we prove Theorem 1, we first show the following lemma .

Lemma 3. *If X and Y have the same homotopy type, then there is a one-to-one correspondence $\psi:\mathcal{A}_\sharp^n(X)\to\mathcal{A}_\sharp^n(Y)$ that is a monoid homomorphism for each nonnegative integer n. Moreover, the restriction of ψ to $\mathcal{E}(X)$ is a group homomorphism into $\mathcal{E}(Y)$.*

Proof. Let $\alpha:X\to Y$ be a homotopy equivalence with homotopy inverse β. Define $\psi:\mathcal{A}_\sharp^n(X)\to\mathcal{A}_\sharp^n(Y)$ by $\psi(f)=\alpha\circ f\circ\beta$ for each $f\in\mathcal{A}_\sharp^n(X)$.

Then, ψ is well-defined, because $f_1\simeq f_2\Rightarrow\alpha\circ f_1\circ\beta\simeq\alpha\circ f_2\circ\beta$ and $\pi_i(\alpha\circ f\circ\beta)=\pi_i(\alpha)\circ\pi_i(f)\circ\pi_i(\beta)$ is an isomorphism on $\pi_i(Y)$ for $i=0,\cdots,n$.

If $\psi(f_1) = \psi(f_2)$, then $\alpha \circ f_1 \circ \beta \simeq \alpha \circ f_2 \circ \beta$. This implies $f_1 \simeq f_2$. Thus, ψ is one-to-one.

For $h \in \mathcal{A}^n_\sharp(Y)$, $\beta \circ h \circ \alpha \in \mathcal{A}^n_\sharp(X)$ and $\psi(\beta \circ h \circ \alpha) = \alpha \circ (\beta \circ h \circ \alpha) \circ \beta \simeq h$. Thus, ψ is onto.

Since

$$\psi(f \circ g) = \alpha \circ (f \circ g) \circ \beta \simeq \alpha \circ (f \circ \beta \circ \alpha \circ g) \circ \beta \simeq (\alpha \circ f \circ \beta) \circ (\alpha \circ g \circ \beta) = \psi(f) \circ \psi(g),$$

ψ is a monoid homomorphism.

If $f \in \mathcal{E}(X)$, then $\alpha \circ f \circ \beta \in \mathcal{E}(Y)$. Thus, $\psi : \mathcal{E}(X) \to \mathcal{E}(Y)$ is an isomorphism.

□

Proof of Theorem 1. It is sufficient to show that $\mathcal{E}(X) = \mathcal{A}^n_\sharp(X)$ if and only if $\mathcal{E}(Y) = \mathcal{A}^n_\sharp(Y)$ for each nonnegative integer n. As $\mathcal{E}(X) \subseteq \mathcal{A}^n_\sharp(X)$ and $\mathcal{E}(Y) \subseteq \mathcal{A}^n_\sharp(Y)$, we will show $\mathcal{A}^n_\sharp(X) \subseteq \mathcal{E}(X)$ if and only if $\mathcal{A}^n_\sharp(Y) \subseteq \mathcal{E}(Y)$. Suppose $\mathcal{A}^n_\sharp(X) \subseteq \mathcal{E}(X)$ and $\psi : \mathcal{A}^n_\sharp(X) \to \mathcal{A}^n_\sharp(Y)$ is the one-to-one correspondence in Lemma 3. If $f \in \mathcal{A}^n_\sharp(Y)$, then $\psi^{-1}(f) \in \mathcal{A}^n_\sharp(X) \subseteq \mathcal{E}(X)$. By Lemma 3, $\psi(\psi^{-1}(f)) \in \mathcal{E}(Y)$. Thus, $f \in \mathcal{E}(Y)$. Consequently, $\mathcal{A}^n_\sharp(Y) \subseteq \mathcal{E}(Y)$.

Similarly, if $\mathcal{A}^n_\sharp(Y) \subseteq \mathcal{E}(Y)$, then $\mathcal{A}^n_\sharp(X) \subseteq \mathcal{E}(X)$.

□

In Lemma 2, we showed that $M(\mathbb{Z}_4, 5)$ is 4-connected whereas its self-closeness number is 5. This is true in general.

Lemma 4. *If X is n-connected and $\mathcal{E}(X) \neq [X, X]$, then $N\mathcal{E}(X) \geq n + 1$.*

Proof. Assume $N\mathcal{E}(X) = k \leq n$. Then, $\mathcal{E}(X) = \mathcal{A}^k_\sharp(X)$ for $k \leq n$. However, because $\pi_i(X) = 0$ for $i = 0, \cdots, n$,

$$\mathcal{A}^n_\sharp(X) = \cdots = \mathcal{A}^k_\sharp(X) = \cdots = \mathcal{A}^0_\sharp(X) = [X, X].$$

This contradicts the hypothesis. Thus, $N\mathcal{E}(X) \geq n + 1$.

□

Corollary 1. *Let X be a CW-complex with cells of dimension $\geq n$, except 0-cells. Then, $N\mathcal{E}(X) \geq n$ provided that $\mathcal{E}(X) \neq [X, X]$.*

The self-closeness number of a CW-complex is closely related to the dimension of the space.

Theorem 2. *If X is a CW-complex with dimension n, then $N\mathcal{E}(X) \leq n$.*

Proof. It is sufficient to show that $\mathcal{A}^n_\sharp(X) = \mathcal{E}(X)$. Let $f \in \mathcal{A}^n_\sharp(X)$. Then $\pi_i(f) \in Aut(\pi_i(X))$ for $i = 0, 1, \cdots, n$. By Theorem 5.1.32 in [2], the induced map $f_\sharp : [X, X] \to [X, X]$ is onto. Thus, there is a map $g : X \to X$ such that $f_\sharp(g) = id_X$, that is, $f \circ g \simeq id_X$. Because $\pi_i(f) \circ \pi_i(g) = \pi_i(f \circ g) = \pi_i(id_X) = id_{\pi_i(X)}$ and $\pi_i(f)$ is an isomorphism, $\pi_i(g) \in Aut(\pi_i(X))$ for $i = 0, 1, \cdots, n$. Thus, $g_\sharp : [X, X] \to [X, X]$ is also onto. Therefore, there is a map $f' : X \to X$ such that $g_\sharp(f') = id_X$, that is, $g \circ f' \simeq id_X$. However, as $f \circ (g \circ f') \simeq f \circ id_X \simeq f$

and $f \circ g \simeq id_X$, $f' \simeq f$. Thus $g \circ f \simeq g \circ f' \simeq id_X$. Therefore, g is a homotopy inverse of f, which implies that $f \in \mathcal{E}(X)$. □

Corollary 2. *Let S^k be the k-dimensional sphere. Then, $N\mathcal{E}(S^k) = k$.*

Proof. As $dim(S^k) = k$, $N\mathcal{E}(S^k) \leq k$ by Theorem 2. It is well known that

$$\mathcal{E}(S^k) \cong \mathbb{Z}_2 \neq \mathbb{Z} \cong [S^k, S^k].$$

However, because S^k is $(k-1)$-connected, $N\mathcal{E}(S^k) \geq k$ by Lemma 4. Thus, $N\mathcal{E}(S^k) = k$. □

In the above corollary, the self-closeness number of a sphere is just its dimension. However, this is not true in general. Before we give an example, we discuss the self-closeness numbers of product spaces.

Theorem 3. *Let X and Y be CW-complexes. Then, we have*

$$N\mathcal{E}(X \times Y) \geq max\{N\mathcal{E}(X), N\mathcal{E}(Y)\}.$$

Proof. Assume that $\mathcal{E}(X \times Y) = \mathcal{A}_\sharp^k(X \times Y)$ for the nonnegative integer $k < max\{N\mathcal{E}(X), N\mathcal{E}(Y)\}$. We may assume that $max\{N\mathcal{E}(X), N\mathcal{E}(Y)\} = N\mathcal{E}(X)$. Let $f \in \mathcal{A}_\sharp^k(X)$ and $g \in \mathcal{A}_\sharp^k(Y)$. Then, $\pi_i(f) \in Aut(\pi_i(X))$ and $\pi_i(g) \in Aut(\pi_i(Y))$ for $i = 0, \cdots, k$. Thus, $\pi_i(f) \times \pi_i(g) \in Aut(\pi_i(X)) \times Aut(\pi_i(Y))$. Let $F : \pi_i(X \times Y) \to \pi_i(X) \times \pi_i(Y)$ be an isomorphism given by $F[r] = ([p_1 r], [p_2 r])$ for $r : S^i \to X \times Y$, where p_i is the projection to the i-th factor for $i = 1, 2$. Then, we have the following commutative diagram:

$$\begin{array}{ccc} \pi_i(X \times Y) & \xrightarrow{\pi_i(f \times g)} & \pi_i(X \times Y) \\ \downarrow F & & \downarrow F \\ \pi_i(X) \times \pi_i(Y) & \xrightarrow{\pi_i(f) \times \pi_i(g)} & \pi_i(X) \times \pi_i(Y). \end{array}$$

It follows that $\pi_i(f \times g)$ is an isomorphism for $i = 0, \cdots, k$. Thus, $f \times g \in \mathcal{A}_\sharp^k(X \times Y) = \mathcal{E}(X \times Y)$. Therefore, $\pi_i(f \times g)$ is an isomorphism for each nonnegative integer i, and so $\pi_i(f) \times \pi_i(g)$ is also an isomorphism for each nonnegative integer i. Thus, $\pi_i(f)$ and $\pi_i(g)$ are isomorphisms for each nonnegative integer i. Consequently, $f \in \mathcal{E}(X)$ and $g \in \mathcal{E}(Y)$. As a result, $\mathcal{E}(X) = \mathcal{A}_\sharp^k(X)$ and $\mathcal{E}(Y) = \mathcal{A}_\sharp^k(Y)$. This contradicts the minimality of $N\mathcal{E}(X)$.

□

Example 3. By the results in [17], $[S^1 \times S^3, S^1 \times S^3] \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_2$ and $\mathcal{E}(S^1 \times S^3) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Thus, by the above theorem,

$$N\mathcal{E}(S^1 \times S^3) \geq max\{N\mathcal{E}(S^1), N\mathcal{E}(S^3)\} = 3.$$

However, as $dim(S^1 \times S^3) = 4$, $N\mathcal{E}(S^1 \times S^3) \leq 4$. Therefore, we have $N\mathcal{E}(S^1 \times S^3) = 3$ or 4. If we give a more detailed computation, we conclude that $N\mathcal{E}(S^1 \times S^3) = 3$. As the dimension of the space is 4, the self-closeness number is not equal to the dimension.

3. The self-closeness number of a Moore space

In this section, we determine the self-closeness numbers of general Moore spaces.

Given an abelian group G and an integer $n \geq 3$, let $M(G,n)$ be a Moore space. Then we obtain the following theorem.

Theorem 4. *$\mathcal{A}_\sharp^n(M(G,n)) = \mathcal{E}(M(G,n))$.*

Proof. Since $\mathcal{E}(M(G,n)) \subseteq \mathcal{A}_\sharp^n(M(G,n))$ in general, it is sufficient to show that $\mathcal{A}_\sharp^n(M(G,n)) \subseteq \mathcal{E}(M(G,n))$. Since $M(G,n)$ is $(n-1)$-connected space, the Hurewicz homomorphism $h_n : \pi_n(M(G,n)) \to H_n(M(G,n))$ is an isomorphism. For any $f \in [M(G,n), M(G,n)]$, the diagram

$$\begin{array}{ccc} \pi_n(M(G,n)) & \xrightarrow{\pi_n(f)} & \pi_n(M(G,n)) \\ {\scriptstyle h_n}\downarrow & & \downarrow{\scriptstyle h_n} \\ H_n(M(G,n)) & \xrightarrow[H_n(f)]{} & H_n(M(G,n)) \end{array}$$

commutes. Let f be an element of $\mathcal{A}_\sharp^n(M(G,n))$. Then $\pi_i(f)$ is an isomorphism for $i \leq n$. Since h_n is an isomorphism, $H_n(f)$ is an isomorphism. Since the Moore space has nontrivial homology only for dimension n, f is a homotopy equivalence by the homology version of the Whitehead Theorem. Hence $\mathcal{A}_\sharp^n(M(G,n)) \subseteq \mathcal{E}(M(G,n))$. □

Since $M(G,n)$ is $(n-1)$-connected space, $\mathcal{A}_\sharp^i(M(G,n)) = [M(G,n), M(G,n)]$ for $i < n$. Moreover, since $[M(G,n), M(G,n)] \neq \mathcal{E}(M(G,n))$, n is the minimum number in the set $\{k \mid \mathcal{E}(M(G,n)) = \mathcal{A}_\sharp^k(M(G,n))\}$. Thus, we have the following corollary.

Corollary 3. *$N\mathcal{E}(M(G,n)) = n$ for $n \geq 3$.*

In [15], Rutter showed that $\mathcal{E}(M(\mathbb{Z}_q,n)) = \mathbb{Z}_{(2,q)} \times \mathbb{Z}_q^*$ where $\mathbb{Z}_q^*$ is the automorphism group of $\mathbb{Z}_q$ and $(2,q)$ is the greatest common divisor of 2 and q. From Theorem 4, $\mathcal{A}_\sharp^n(M(\mathbb{Z}_q,n)) = \mathcal{E}(M(\mathbb{Z}_q,n))$ for all $q > 1$. Moreover, since $\mathcal{E}(M(\mathbb{Z}_q,n)) \subset \mathcal{A}_\sharp^k(M(\mathbb{Z}_q,n)) \subset \mathcal{A}_\sharp^n(M(\mathbb{Z}_q,n))$ if $n \leq k$, we have

$$\mathcal{A}_\sharp^k(M(\mathbb{Z}_q,n)) = \begin{cases} [M(\mathbb{Z}_q,n), M(\mathbb{Z}_q,n)] & \text{if } k < n, \\ \mathbb{Z}_q^* & \text{if } k \geq n,\ q \text{ is odd}, \\ \mathbb{Z}_2 \times \mathbb{Z}_q^* & \text{if } k \geq n,\ q \text{ is even}. \end{cases}$$

4. An equivalence relation on $\mathcal{A}_\sharp^n(X)$

We define a relation '$\simeq_n$' on $\mathcal{A}_\sharp^n(X)$ as follows: for $f, g \in \mathcal{A}_\sharp^n(X)$, $f \simeq_n g$ if $\pi_i(f) = \pi_i(g) : \pi_i(X) \to \pi_i(X)$ for $i = 0, \cdots, n$. If $f \simeq_n g$, then we say *f is n-self homotopic to g*. By definition, '$\simeq_n$' is an equivalence relation. We denote by $\bar{f}$ the equivalence class of f in $\mathcal{A}_\sharp^n(X)$, and call this the *n-self homotopy*

equivalence class of f on X. Moreover, we denote by $\bar{\mathcal{A}}_\sharp^n(X)$ the set of all n-self homotopy equivalence classes on X. $n = \infty$ is available, and we denote $\bar{\mathcal{A}}_\sharp^\infty(X)$ as simply $\bar{\mathcal{A}}_\sharp(X)$.

In general, $\bar{\mathcal{A}}_\sharp^n(X) \neq \mathcal{A}_\sharp^n(X)$.

Example 4. By [5], It was shown that $\mathcal{E}_\sharp(S^1 \times S^3) = \mathbb{Z}_2$. However, $\mathcal{A}_\sharp(S^1 \times S^3) = \mathcal{E}(S^1 \times S^3) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Choose two distinct elements $\alpha, \beta \in \mathcal{E}_\sharp(S^1 \times S^3) = \mathbb{Z}_2$. Then since $\pi_i(\alpha) = id_{\pi_i(X)} = \pi_i(\beta)$, $\bar{\alpha} = \bar{\beta}$ in $\bar{\mathcal{A}}_\sharp(S^1 \times S^3)$.

Theorem 5. *Let X be a CW-complex. Then, $\bar{\mathcal{A}}_\sharp^n(X)$ is a monoid for $n < \infty$, and $\bar{\mathcal{A}}_\sharp(X)$ is a group.*

Proof. Let us define $\bar{f} \cdot \bar{g} = \overline{f \circ g}$ for $f, g \in \bar{\mathcal{A}}_\sharp^n(X)$. Then, the operation '$\cdot$' is well-defined and associative. Furthermore, $\overline{id_X}$ is an element of $\bar{\mathcal{A}}_\sharp^n(X)$. Thus, $\bar{\mathcal{A}}_\sharp^n(X)$ has a monoid structure.

As $\mathcal{A}_\sharp(X) = \mathcal{E}(X)$ by Lemma 2, for each $\bar{f} \in \mathcal{A}_\sharp(X)$, its representative f belongs to $\mathcal{E}(X)$. Thus, f has the homotopy inverse g in $\mathcal{E}(X) = \mathcal{A}_\sharp(X)$. Then, $\bar{g}$ is the inverse element of $\bar{f}$. As a result, $\bar{\mathcal{A}}_\sharp(X)$ has a group structure. □

Corollary 4. *Let X be a CW-complex. If $N\mathcal{E}(X) = n$, then $\bar{\mathcal{A}}_\sharp^k(X)$ has a group structure for $k \geq n$.*

Consider the n-self equivalence class $\bar{f}$ of $f \in \mathcal{A}_\sharp^n(X)$. Then, $\bar{f}$ is a subset of $\mathcal{A}_\sharp^n(X)$. In particular, $\overline{id_X}$ is a submonoid of $\mathcal{A}_\sharp^n(X)$. For a given $g \in \mathcal{A}_\sharp^n(X)$, we define a set $g \cdot \bar{f}$ as follows:

$$g \cdot \bar{f} := \{g \circ \alpha | \alpha \in \bar{f}\}.$$

Similarly, we define the set $\bar{f} \cdot g$ as

$$\bar{f} \cdot g := \{\beta \circ g | \beta \in \bar{f}\}.$$

In general, $g \cdot \bar{f} \subseteq \overline{g \circ f}$, $\bar{f} \cdot g \subseteq \overline{f \circ g}$, and $g \cdot \bar{f} \neq \bar{f} \cdot g$. Moreover, $\mathcal{E}_\sharp^n(X) \subseteq \overline{id_X} \subseteq \mathcal{A}_\sharp^n(X)$ for each positive integer n. In fact, if $f \in \mathcal{E}_\sharp^n(X)$, then $\pi_i(f) = id_{\pi_i(X)}$ for $i = 0, \cdots, n$. However, $\pi_i(f) = id_{\pi_i(X)} = \pi_i(id_X)$. Thus, $f \in \overline{id_X}$.

Lemma 5. *$\mathcal{E}_\sharp^n(X)$ is a normal subgroup of $\mathcal{E}(X)$ for all n.*

Proof. Let $h \in \mathcal{E}_\sharp^n(X)$ and $f \in \mathcal{E}(X)$. Then, for the homotopy inverse g of f, we have

$$\pi_i(f \circ h \circ g) = \pi_i(f) \circ \pi_i(h) \circ \pi_i(g) = \pi_i(f) \circ \pi_i(g) = \pi_i(id_X) = id_{\pi_i(X)}$$

for $i = 0, \cdots, n$. Thus, $f \circ h \circ g \in \mathcal{E}_\sharp^n(X)$. □

By the above lemma, we can consider the factor group $\mathcal{E}(X)/\mathcal{E}_\sharp^n(X)$. Here we investigate the relation between this factor group and the n-equivalence classes.

Theorem 6. *$\mathcal{E}(X)/\mathcal{E}^n_\sharp(X)$ is isomorphic to a subgroup of $\bar{\mathcal{A}}^n_\sharp(X)$ for each positive integer n. In particular, if $N\mathcal{E}(X) = n$, then $\bar{\mathcal{A}}^n_\sharp(X) \cong \mathcal{E}(X)/\mathcal{E}^n_\sharp(X)$.*

Proof. Define $\Psi : \mathcal{E}(X)/\mathcal{E}^n_\sharp(X) \to \bar{\mathcal{A}}^n_\sharp(X)$ by $\Psi(f \cdot \mathcal{E}^n_\sharp(X)) = \bar{f}$. If $f \cdot \mathcal{E}^n_\sharp(X) = g \cdot \mathcal{E}^n_\sharp(X)$ for $f, g \in \mathcal{E}(X)$, then $g^{-1} \circ f \in \mathcal{E}^n_\sharp(X)$. Thus, $id_{\pi_i(X)} = \pi_i(g^{-1} \circ f) = \pi_i(g^{-1}) \circ \pi_i(f)$ for $i = 0, \cdots, n$. It follows that $\pi_i(g) = \pi_i(f)$ for $i = 0, \cdots, n$ and $\bar{f} = \bar{g}$. Thus, we conclude that Ψ is well-defined.

If $\Psi(f \cdot \mathcal{E}^n_\sharp(X)) = \Psi(g \cdot \mathcal{E}^n_\sharp(X))$, then $\bar{f} = \bar{g}$ in $\bar{\mathcal{A}}^n_\sharp(X)$. Thus, $\pi_i(g) = \pi_i(f)$ for $i = 0, \cdots, n$. Therefore, $id_{\pi_i(X)} = \pi_i(g)^{-1} \circ \pi_i(f) = \pi_i(g^{-1} \circ f)$ for $i = 0, \cdots, n$. It follows that $g^{-1} \circ f \in \mathcal{E}^n_\sharp(X)$. That is, $f \cdot \mathcal{E}^n_\sharp(X) = g \cdot \mathcal{E}^n_\sharp(X)$. Thus, Ψ is one-to-one.

Furthermore, since $\Psi((f \cdot \mathcal{E}^n_\sharp(X)) \cdot (g \cdot \mathcal{E}^n_\sharp(X))) = (f \circ g) \cdot \mathcal{E}^n_\sharp(X) = \overline{f \circ g} = \bar{f} \cdot \bar{g} = \Psi((f \cdot \mathcal{E}^n_\sharp(X)) \cdot \Psi((g \cdot \mathcal{E}^n_\sharp(X))$, Ψ is a monomorphism.

In the case that $N\mathcal{E}(X) = n$, Ψ is onto. In fact, if $\bar{f} \in \bar{\mathcal{A}}^n_\sharp(X)$, then $f \in \mathcal{A}^n_\sharp(X) = \mathcal{E}(X)$ and $\Psi(f \cdot \mathcal{E}^n_\sharp(X)) = \bar{f}$.

□

Corollary 5. $\bar{\mathcal{A}}_\sharp(X) \cong \mathcal{E}(X)/\mathcal{E}_\sharp(X)$.

Corollary 6. *If $\mathcal{E}^n_\sharp(X) = \{id_X\}$ for some positive integer n, $\bar{\mathcal{A}}_\sharp(X) \cong \mathcal{E}(X)$.*

Proof. Let $\mathcal{E}^n_\sharp(X) = \{id_X\}$. Since $\mathcal{E}_\sharp(X) \subseteq \cdots \subseteq \mathcal{E}^n_\sharp(X)$, $\mathcal{E}_\sharp(X) = \{id_X\}$. Thus, $\bar{\mathcal{A}}_\sharp(X) \cong \mathcal{E}(X)$ by Corollary 5. □

Example 5. In [8], It was shown that $\mathcal{E}^{n+2}_\sharp(M(\mathbb{Z}_q, n+1) \vee (M(\mathbb{Z}_p, n)) = \{id_X\}$ for $n > 4$ if q and p are odd. Thus,

$$\bar{\mathcal{A}}_\sharp(M(\mathbb{Z}_q, n+1) \vee (M(\mathbb{Z}_p, n)) \cong \mathcal{E}(M(\mathbb{Z}_q, n+1) \vee (M(\mathbb{Z}_p, n))$$

by Corollary 6.

Acknowledgements

We would like to thank the referee(s) for offering other suggestions and comments that improved the quality of the paper.

References

[1] M. Arkowitz, *The group of self-homotopy equivalences a survey*, Lecture Notes in Math. 1425 Springer, New York, (1990), 170-203.

[2] M. Aquilar, S. Gitler and C. Prieto, *Algebraic topology from a homotopical viewpoint*, Springer-Verlag New York, Inc. (2000).

[3] M. Arkowitz and Lupton, *On finiteness of subgroups of self-homotopy equivalences*, Contemp. Math. 181 Amer. Math. Soc., (1995), 1-25.

[4] M. Arkowitz and K. Maruyama, *Self-homotopy equivalences which induce the identity on homology, cohomology or homotopy groups*, Topology Appl. 87 no. 2, (1998), 133-154.

[5] M. Arkowitz, H. Oshima and J. Strom, *Homotopy classes of self-maps and induced homomorphisms of homotopy groups*, J. Math. Soc. Japan 58, no. 2 (2006), 401-418.

[6] S. Araki and Toda, *Multiplicative structure in mod q cohomology theories I*, Osaka J. Math. 2 (1965), 71-115.
[7] A. L. Blackers and W. S. Massey, *The homotopy groups of triads II*, Ann. of Math. no 55, (1952), 192-201.
[8] H. Choi and K. Lee, *Certain self homotopy equivalences on wedge product on Moore spaces*, to appear.
[9] B. Gray. *Homotopy Theory*, Academic Press, Inc., (1975).
[10] P. Hilton, *Homotopy Theory and Duality*, Gordon and Breach, New York, 1965.
[11] M. Jeong, *Certain subgroups of self homotopy equivalences of the wedge of two Moore spaces*, Commun. Korean Math. Soc. 25, no. 1 (2010), 111-117.
[12] K. Lee, *The groups of self pair homotopy equivalences*, Journal of Korea Math. Soc. 43, no. 3, (2006), 491-506.
[13] S. Oka, *On the subgroup of self-equivalences of H-space of low rank*, 1,2, Mem. Fac. Sci. Kyushu Univ. Ser A 35, no. 2, (1981), 247-282.
[14] D. Puppe, *homotopiemenge und ihre induzierten Abbildungen*, I, Math. Z. no 69, (1958), 425-505.
[15] J. Rutter, *The group of self-equivalence classes of CW-complexes*, Math. Proc. Cambridge Phil. Soc. 93 , no. 2,(1983), 275-293.
[16] N. Sawashita, *On the group of self-equivalences of the product spheres*, Hiroshima Math. J. 5, (1975), 69-86.
[17] A. Sieradski, *Twisted self-homotopy equivalences*, Pacific J. Math. 34, (1970), 789-802.
[18] E. Spanier, *Algebraic topology*, McGraw-Hill, New York, (1966).
[19] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies, no 49, (1962).

Department of Mathematics
Korea University
Seoul 702-701, Korea
E-mail address: howon@korea.ac.kr

Department of Information and Mathematics
Korea University
Sejong 425-791, Korea
E-mail address: keyolee@korea.ac.kr